

# Fuzzy $\alpha^m$ -Separation Axioms

<sup>1</sup>Qays Hatem Imran, <sup>2</sup>Hayder Kadhim Zghair

<sup>1</sup>Department of Mathematics, College of Education for Pure Science, Al-Muthanna University, Samawah, Iraq

<sup>2</sup>Department of Software, College of Information Technology, University of Babylon, Babylon, Iraq

**Abstract:** This paper will introduce a new class of fuzzy closed set (briefly F-CS) called fuzzy  $\alpha^m$ -closed set as well as, introduce the fuzzy  $\alpha^m$ -kernel set of the fuzzy topological space. The investigation will address and discuss some of the properties of the fuzzy separation axioms such as fuzzy  $\alpha^m$ - $R_i$ -space and fuzzy  $\alpha^m$ - $T_j$ -space (note that, the indexes  $i$  and  $j$  are natural numbers of the spaces  $R$  and  $T$  are from 0 to 3 and from 0 to 4 respectively).

**Mathematics Subject Classification (2010):** 54A40.

**Keywords:** Fuzzy  $\alpha^m$ -closed set, Fuzzy  $\alpha^m$ - $R_i$ -space,  $i = 0, 1, 2, 3$  and Fuzzy  $\alpha^m$ - $T_j$ -space,  $j = 0, 1, 2, 3, 4$ .

## 1. INTRODUCTION

In 1965 Zadeh studied the fuzzy sets (briefly F-sets) (see [6]) which plays such a role in the field of fuzzy topological spaces (or simply fts). The fuzzy topological spaces investigated by Chang in 1968 (see [3]). A. S. Bin Shahna [1] defined fuzzy  $\alpha$ -closed sets. In 1997, fuzzy generalized closed set (briefly Fg-CS) was introduced by G. Balasubramania and P. Sundaram [5]. In 2014, M. Mathew and R. Parimelazhagan [7] defined  $\alpha^m$ -closed sets of topological spaces. The aim of this paper is to introduce a concept of Fuzzy  $\alpha^m$ -closed sets and study their basic properties in fts. Furthermore, the investigation will include some of the properties of the fuzzy separation axioms such as fuzzy  $\alpha^m$ - $R_i$ -space and fuzzy  $\alpha^m$ - $T_j$ -space (here the indexes  $i$  and  $j$  are natural numbers of the spaces  $R$  and  $T$  are from 0 to 3 and from 0 to 4 respectively).

## 2. PRELIMINARIES

Throughout this paper,  $(X, \tau)$  or simply  $X$  always mean a fts. A fuzzy point [4] with support  $x \in X$  and value  $\lambda$  ( $0 < \lambda \leq 1$ ) at  $x \in X$  will be denoted by  $x_\lambda$ , and for fuzzy set  $\mathcal{A}$ ,  $x_\lambda \in \mathcal{A}$  iff  $\lambda \leq \mathcal{A}(x)$ . Two fuzzy points  $x_\lambda$  and  $y_\sigma$  are said to be distinct iff their supports are distinct. That is, by  $0_X$  and  $1_X$  we mean the constant fuzzy sets taking the values 0 and 1 on  $X$ , respectively [2]. For a fuzzy set  $\mathcal{A}$  in a fts  $(X, \tau)$ ,  $cl(\mathcal{A})$ ,  $int(\mathcal{A})$  and  $\mathcal{A}^c = 1_X - \mathcal{A}$  represents the fuzzy closure of  $\mathcal{A}$ , the fuzzy interior of  $\mathcal{A}$  and the fuzzy complement of  $\mathcal{A}$  respectively.

**Definition 2.1:[12]** A fuzzy point in a set  $X$  with support  $x$  and membership value 1 is called crisp point, denoted by  $x_1$ . For any fuzzy set  $\mathcal{A}$  in  $X$ , we have  $x_1 \in \mathcal{A}$  iff  $\mathcal{A}(x) = 1$ .

**Definition 2.2:[8]** A fuzzy point  $x_\lambda \in \mathcal{A}$  is called quasi-coincident (briefly  $q$ -coincident) with the fuzzy set  $\mathcal{A}$  is denoted by  $x_\lambda q \mathcal{A}$  iff  $\lambda + \mathcal{A}(x) > 1$ . A fuzzy set  $\mathcal{A}$  in a fts  $(X, \tau)$  is called  $q$ -coincident with a fuzzy set  $\mathcal{B}$  which is denoted by  $\mathcal{A} q \mathcal{B}$  iff there exists  $x \in X$  such that  $\mathcal{A}(x) + \mathcal{B}(x) > 1$ . If the fuzzy sets  $\mathcal{A}$  and  $\mathcal{B}$  in a fts  $(X, \tau)$  are not  $q$ -coincident then we write  $\mathcal{A} \bar{q} \mathcal{B}$ . Note that  $\mathcal{A} \leq \mathcal{B} \Leftrightarrow \mathcal{A} \bar{q} (1_X - \mathcal{B})$ .

**Definition 2.3:[8]** A fuzzy set  $\mathcal{A}$  in a fts  $(X, \tau)$  is called  $q$ -neighbourhood (briefly  $q$ -nhd) of a fuzzy point  $x_\lambda$  (resp. fuzzy set  $\mathcal{B}$ ) if there is a F-OS  $\mathcal{M}$  in a fts  $(X, \tau)$  such that  $x_\lambda q \mathcal{M} \leq \mathcal{A}$  (resp.  $\mathcal{B} q \mathcal{M} \leq \mathcal{A}$ ).

**Definition 2.4:[1]** A fuzzy set  $\mathcal{A}$  of a fts  $(X, \tau)$  is called a fuzzy  $\alpha$ -open set (briefly  $F\alpha$ -OS) if  $\mathcal{A} \leq \text{int}(cl(\text{int}(\mathcal{A})))$  and a fuzzy  $\alpha$ -closed set (briefly  $F\alpha$ -CS) if  $cl(\text{int}(cl(\mathcal{A}))) \leq \mathcal{A}$ . The fuzzy  $\alpha$ -closure of a fuzzy set  $\mathcal{A}$  of fts  $(X, \tau)$  is the intersection of all  $F\alpha$ -CS that contain  $\mathcal{A}$  and is denoted by  $\alpha cl(\mathcal{A})$ .

**Definition 2.5:[5]** A fuzzy set  $\mathcal{A}$  of a fts  $(X, \tau)$  is called a fuzzy g-closed set (briefly Fg-CS) if  $cl(\mathcal{A}) \leq \mathcal{U}$  whenever  $\mathcal{A} \leq \mathcal{U}$  and  $\mathcal{U}$  is a F-OS in  $X$ .

**Definition 2.6:[10]** A fuzzy set  $\mathcal{A}$  of a fts  $(X, \tau)$  is called a fuzzy  $\alpha g$ -closed set (briefly  $F\alpha g$ -CS) if  $\alpha cl(\mathcal{A}) \leq \mathcal{U}$  whenever  $\mathcal{A} \leq \mathcal{U}$  and  $\mathcal{U}$  is a  $F\alpha$ -OS in  $X$ .

**Definition 2.7:[9]** A fuzzy set  $\mathcal{A}$  of a fts  $(X, \tau)$  is called a fuzzy  $g\alpha$ -closed set (briefly  $Fg\alpha$ -CS) if  $\alpha cl(\mathcal{A}) \leq \mathcal{U}$  whenever  $\mathcal{A} \leq \mathcal{U}$  and  $\mathcal{U}$  is a F-OS in  $X$ .

**Remark 2.8:[5,11]** In a fts  $(X, \tau)$ , then the following statements are true:

- (i) Every F-CS is a Fg-CS.
- (ii) Every F-CS is a  $F\alpha$ -CS.

**Remark 2.9:[9,10]** In a fts  $(X, \tau)$ , then the following statements are true:

- (i) Every Fg-CS is a  $Fg\alpha$ -CS.
- (ii) Every  $F\alpha$ -CS is a  $F\alpha g$ -CS.
- (iii) Every  $F\alpha g$ -CS is a  $Fg\alpha$ -CS.

### 3. FUZZY $\alpha^m$ -CLOSED SETS

**Definition 3.1:** A fuzzy set  $\mathcal{A}$  of a fts  $(X, \tau)$  is called a fuzzy  $\alpha^m$ -closed set (briefly  $F\alpha^m$ -CS) if  $\text{int}(cl(\mathcal{A})) \leq \mathcal{U}$  whenever  $\mathcal{A} \leq \mathcal{U}$  and  $\mathcal{U}$  is a  $F\alpha$ -OS. The complement of a fuzzy  $\alpha^m$ -closed set in  $X$  is fuzzy  $\alpha^m$ -open set (briefly  $F\alpha^m$ -OS) in  $X$ , the family of all  $F\alpha^m$ -OS (resp.  $F\alpha^m$ -CS) of a fts  $(X, \tau)$  is denoted by  $F\alpha^m\text{-O}(X)$  (resp.  $F\alpha^m\text{-C}(X)$ ).

**Example 3.2:** Let  $X = \{x, y\}$  and the fuzzy set  $\mathcal{A}$  in  $X$  defined as follows:  $\mathcal{A}(x) = 0.5, \mathcal{A}(y) = 0.5$ . Let  $\tau = \{0_X, \mathcal{A}, 1_X\}$  be a fts. Then the fuzzy sets  $0_X, \mathcal{A}$  and  $1_X$  are  $F\alpha^m$ -OS and  $F\alpha^m$ -CS at the same time in  $X$ .

**Remark 3.3:** In a fts  $(X, \tau)$ , then the following statements are true:

- (i) Every F-CS is a  $F\alpha^m$ -CS.
- (ii) Every  $F\alpha^m$ -CS is a  $F\alpha$ -CS.
- (iii) Every  $F\alpha^m$ -CS is a  $F\alpha g$ -CS.
- (iv) Every  $F\alpha^m$ -CS is a  $Fg\alpha$ -CS.

**Proof:** (i) This follows directly from the definition (3.1).

(ii) Let  $\mathcal{A}$  be a  $F\alpha^m$ -CS in  $X$  and let  $\mathcal{U}$  be a F-OS such that  $\mathcal{A} \leq \mathcal{U}$ . Since every F-OS is a  $F\alpha$ -OS and  $\mathcal{A}$  is a  $F\alpha^m$ -CS,  $\text{int}(cl(\mathcal{A})) \leq (\text{int}(cl(\mathcal{A}))) \vee (cl(\text{int}(\mathcal{A}))) \leq \mathcal{U}$ . Therefore,  $\mathcal{A}$  is a  $F\alpha$ -CS in  $X$ .

(iii) From the part (ii) and remark (2.9) (ii).

(iv) From the part (iii) and remark (2.9) (iii).

**Theorem 3.4:** A fuzzy set  $\mathcal{A}$  is  $F\alpha^m$ -CS iff  $\text{int}(cl(\mathcal{A})) - \mathcal{A}$  contains no non-empty  $F\alpha^m$ -CS.

**Proof: Necessity.** Suppose that  $\mathcal{F}$  is a non-empty  $F\alpha^m$ -closed subset of  $\text{int}(cl(\mathcal{A}))$  such that  $\mathcal{F} < \text{int}(cl(\mathcal{A})) - \mathcal{A}$ . Then  $\mathcal{F} \leq \text{int}(cl(\mathcal{A})) - \mathcal{A}$ . Then  $\mathcal{F} \leq \text{int}(cl(\mathcal{A})) \wedge \mathcal{A}^c$ . Therefore  $\mathcal{F} \leq \text{int}(cl(\mathcal{A}))$  and  $\mathcal{F} \leq \mathcal{A}^c$ . Since  $\mathcal{F}^c$  is a  $F\alpha^m$ -OS and  $\mathcal{A}$  is a  $F\alpha^m$ -CS,  $\text{int}(cl(\mathcal{A})) \leq \mathcal{F}^c$ . Thus  $\mathcal{F} \leq (\text{int}(cl(\mathcal{A})))^c$ . Therefore  $\mathcal{F} \leq (\text{int}(cl(\mathcal{A}))) \wedge (\text{int}(cl(\mathcal{A})))^c = 0_X$ . Therefore  $\mathcal{F} = 0_X \Rightarrow \text{int}(cl(\mathcal{A})) - \mathcal{A}$  contains no non-empty  $F\alpha^m$ -CS.

**Sufficiency.** Let  $\mathcal{A} \leq \mathcal{U}$  be a  $F\alpha^m$ -OS. Suppose that  $int(cl(\mathcal{A}))$  is not contained in  $\mathcal{U}$ . Then  $(int(cl(\mathcal{A})))^c$  is a non-empty  $F\alpha^m$ -CS and contained in  $int(cl(\mathcal{A})) - \mathcal{A}$  which is a contradiction. Therefore,  $int(cl(\mathcal{A})) \leq \mathcal{U}$  and hence  $\mathcal{A}$  is a  $F\alpha^m$ -CS.

**Theorem 3.5:** Let  $\mathcal{B} \leq Y \leq X$ , if  $\mathcal{B}$  is a  $F\alpha^m$ -CS relative to  $Y$  and  $Y$  is a F-OS then  $\mathcal{B}$  is a  $F\alpha^m$ -CS in a fts  $(X, \tau)$ .

**Proof:** Let  $\mathcal{U}$  be a  $F\alpha$ -OS in a fts  $(X, \tau)$  such that  $\mathcal{B} \leq \mathcal{U}$ . Given that  $\mathcal{B} \leq Y \leq X$ . Therefore  $\mathcal{B} \leq Y$  and  $\mathcal{B} \leq \mathcal{U}$ . This implies  $\mathcal{B} \leq Y \wedge \mathcal{U}$ . Since  $\mathcal{B}$  is a  $F\alpha^m$ -CS relative to  $Y$ , then  $int(cl(\mathcal{B})) \leq \mathcal{U}$ .  $Y \wedge int(cl(\mathcal{B})) \leq Y \wedge \mathcal{U}$  implies that  $Y \wedge (int(cl(\mathcal{B}))) \leq \mathcal{U}$ . Thus  $[Y \wedge int(cl(\mathcal{B}))] \vee [int(cl(\mathcal{B}))]^c \leq \mathcal{U} \vee [int(cl(\mathcal{B}))]^c$ . This implies that  $(Y \vee (int(cl(\mathcal{B})))^c) \wedge (int(cl(\mathcal{B}))) \vee (int(cl(\mathcal{B})))^c \leq \mathcal{U} \vee (int(cl(\mathcal{B})))^c$ . Therefore  $(Y < (int(cl(\mathcal{B})))^c) \leq \mathcal{U} \vee (int(cl(\mathcal{B})))^c$ . Since  $Y$  is a F-OS in  $X$ ,  $int(cl(Y)) \leq \mathcal{U} \vee (int(cl(\mathcal{B})))^c$ . Also  $\mathcal{B} \leq Y$  implies that  $int(cl(\mathcal{B})) \leq int(cl(Y))$ . Thus  $int(cl(\mathcal{B})) \leq int(cl(Y)) \leq \mathcal{U} \vee (int(cl(\mathcal{B})))^c$ . Therefore  $int(cl(\mathcal{B})) \leq \mathcal{U}$ . Since  $int(cl(\mathcal{B}))$  is not contained in  $(int(cl(\mathcal{B})))^c$ ,  $\mathcal{B}$  is a  $F\alpha^m$ -CS relative to  $X$ .

**Theorem 3.6:** If  $\mathcal{A}$  is a  $F\alpha^m$ -CS and  $\mathcal{A} \leq \mathcal{B} \leq int(cl(\mathcal{A}))$ , then  $\mathcal{B}$  is a  $F\alpha^m$ -CS.

**Proof:** Let  $\mathcal{A}$  be a  $F\alpha^m$ -CS such that  $\mathcal{A} \leq \mathcal{B} \leq int(cl(\mathcal{A}))$ . Let  $\mathcal{U}$  be a  $F\alpha$ -OS in a fts  $(X, \tau)$  such that  $\mathcal{B} \leq \mathcal{U}$ . Since  $\mathcal{A}$  is a  $F\alpha^m$ -CS, we have  $int(cl(\mathcal{A})) \leq \mathcal{U}$  whenever  $\mathcal{A} \leq \mathcal{U}$ . Since  $\mathcal{A} \leq \mathcal{B}$  and  $\mathcal{B} \leq int(cl(\mathcal{A}))$ , then  $int(cl(\mathcal{B})) \leq int(cl(int(cl(\mathcal{A})))) \leq int(cl(\mathcal{A})) \leq \mathcal{U}$ . Therefore,  $int(cl(\mathcal{B})) \leq \mathcal{U}$ . Thus,  $\mathcal{B}$  is a  $F\alpha^m$ -CS in  $X$ .

**Theorem 3.7:** The intersection of a  $F\alpha^m$ -CS and a F-CS is a  $F\alpha^m$ -CS.

**Proof:** Let  $\mathcal{A}$  be a  $F\alpha^m$ -CS and  $\mathcal{F}$  be a F-CS. Since  $\mathcal{A}$  is a  $F\alpha^m$ -CS,  $int(cl(\mathcal{A})) \leq \mathcal{U}$  whenever  $\mathcal{A} \leq \mathcal{U}$  where  $\mathcal{U}$  is a  $F\alpha$ -OS. To show that  $\mathcal{A} \wedge \mathcal{F}$  is a  $F\alpha^m$ -CS. It is enough to show that  $int(cl(\mathcal{A} \wedge \mathcal{F})) \leq \mathcal{U}$  whenever  $\mathcal{A} \wedge \mathcal{F} \leq \mathcal{U}$ , where  $\mathcal{U}$  is a  $F\alpha$ -OS. Let  $\mathcal{M} = 1_X - \mathcal{F}$  then  $\mathcal{A} \leq \mathcal{U} \vee \mathcal{M}$ . Since  $\mathcal{M}$  is a F-OS,  $\mathcal{U} \vee \mathcal{M}$  is a  $F\alpha$ -OS and  $\mathcal{A}$  is a  $F\alpha^m$ -CS,  $int(cl(\mathcal{A})) \leq \mathcal{U} \vee \mathcal{M}$ . Now,  $int(cl(\mathcal{A} \wedge \mathcal{F})) \leq int(cl(\mathcal{A})) \wedge int(cl(\mathcal{F})) \leq int(cl(\mathcal{A})) \wedge \mathcal{F} \leq (\mathcal{U} \vee \mathcal{M}) \wedge \mathcal{F} \leq (\mathcal{U} \wedge \mathcal{F}) \vee (\mathcal{M} \wedge \mathcal{F}) \leq (\mathcal{U} \wedge \mathcal{F}) \vee 0_X \leq \mathcal{U}$ . This implies that  $\mathcal{A} \wedge \mathcal{F}$  is a  $F\alpha^m$ -CS.

**Theorem 3.8:** If  $\mathcal{A}$  and  $\mathcal{B}$  are two  $F\alpha^m$ -CS in a fts  $(X, \tau)$ , then  $\mathcal{A} \wedge \mathcal{B}$  is a  $F\alpha^m$ -CS in  $X$ .

**Proof:** Let  $\mathcal{A}$  and  $\mathcal{B}$  be two  $F\alpha^m$ -CS in a fts  $(X, \tau)$ . Let  $\mathcal{U}$  be a  $F\alpha$ -OS in  $X$  such that  $\mathcal{A} \wedge \mathcal{B} \leq \mathcal{U}$ . Now,  $int(cl(\mathcal{A} \wedge \mathcal{B})) \leq int(cl(\mathcal{A})) \wedge int(cl(\mathcal{B})) \leq \mathcal{U}$ . Hence  $\mathcal{A} \wedge \mathcal{B}$  is a  $F\alpha^m$ -CS.

**Remark 3.9:** The union of two  $F\alpha^m$ -CS need not be a  $F\alpha^m$ -CS.

**Definition 3.10:** The intersection of all  $F\alpha^m$ -CS in a fts  $(X, \tau)$  containing  $\mathcal{A}$  is called fuzzy  $\alpha^m$ -closure of  $\mathcal{A}$  and is denoted by  $\alpha^m-cl(\mathcal{A})$ ,  $\alpha^m-cl(\mathcal{A}) = \bigwedge \{\mathcal{B} : \mathcal{A} \leq \mathcal{B}, \mathcal{B} \text{ is a } F\alpha^m\text{-CS}\}$ .

**Definition 3.11:** The union of all  $F\alpha^m$ -OS in a fts  $(X, \tau)$  contained in  $\mathcal{A}$  is called fuzzy  $\alpha^m$ -interior of  $\mathcal{A}$  and is denoted by  $\alpha^m-int(\mathcal{A})$ ,  $\alpha^m-int(\mathcal{A}) = \bigvee \{\mathcal{B} : \mathcal{B} \geq \mathcal{A}, \mathcal{B} \text{ is a } F\alpha^m\text{-OS}\}$ .

**Proposition 3.12:** Let  $\mathcal{A}$  be any fuzzy set in a fts  $(X, \tau)$ . Then the following properties hold:

- (i)  $\alpha^m-int(\mathcal{A}) = \mathcal{A}$  iff  $\mathcal{A}$  is a  $F\alpha^m$ -OS.
- (ii)  $\alpha^m-cl(\mathcal{A}) = \mathcal{A}$  iff  $\mathcal{A}$  is a  $F\alpha^m$ -CS.
- (iii)  $\alpha^m-int(\mathcal{A})$  is the largest  $F\alpha^m$ -OS contained in  $\mathcal{A}$ .
- (iv)  $\alpha^m-cl(\mathcal{A})$  is the smallest  $F\alpha^m$ -CS containing  $\mathcal{A}$ .

**Proof:** (i), (ii), (iii) and (iv) are obvious.

**Proposition 3.13:** Let  $\mathcal{A}$  be any fuzzy set in a fts  $(X, \tau)$ . Then the following properties hold:

$$(i) \alpha^m\text{-int}(1_X - \mathcal{A}) = 1_X - (\alpha^m\text{-cl}(\mathcal{A})),$$

$$(ii) \alpha^m\text{-cl}(1_X - \mathcal{A}) = 1_X - (\alpha^m\text{-int}(\mathcal{A})).$$

**Proof:** (i) By definition,  $\alpha^m\text{-cl}(\mathcal{A}) = \bigwedge \{\mathcal{B}: \mathcal{A} \leq \mathcal{B}, \mathcal{B} \text{ is a } F\alpha^m\text{-CS}\}$

$$\begin{aligned} 1_X - (\alpha^m\text{-cl}(\mathcal{A})) &= 1_X - \bigwedge \{\mathcal{B}: \mathcal{A} \leq \mathcal{B}, \mathcal{B} \text{ is a } F\alpha^m\text{-CS}\} \\ &= \bigvee \{1_X - \mathcal{B}: \mathcal{A} \leq \mathcal{B}, \mathcal{B} \text{ is a } F\alpha^m\text{-CS}\} \\ &= \bigvee \{\mathcal{H}: 1_X - \mathcal{A} \geq \mathcal{H}, \mathcal{H} \text{ is a } F\alpha^m\text{-OS}\} \\ &= \alpha^m\text{-int}(1_X - \mathcal{A}) \end{aligned}$$

(ii) The proof is similar to (i).

**Definition 3.14:** A fuzzy set  $\mathcal{A}$  in a fts  $(X, \tau)$  is said to be a fuzzy  $\alpha^m$ -neighbourhood (briefly  $F\alpha^m$ -nhd) of a fuzzy point  $x_\lambda$  if there exists a  $F\alpha^m$ -OS  $\mathcal{B}$  such that  $x_\lambda \in \mathcal{B} \leq \mathcal{A}$ . A  $F\alpha^m$ -nhd  $\mathcal{A}$  is said to be a  $F\alpha^m$ -open-nhd (resp.  $F\alpha^m$ -closed-nhd) iff  $\mathcal{A}$  is a  $F\alpha^m$ -OS (resp.  $F\alpha^m$ -CS). A fuzzy set  $\mathcal{A}$  in a fts  $(X, \tau)$  is said to be a fuzzy  $\alpha^m$ - $q$ -neighbourhood (briefly  $F\alpha^m$ - $q$ -nhd) of a fuzzy point  $x_\lambda$  (resp. fuzzy set  $\mathcal{B}$ ) if there exists a  $F\alpha^m$ -OS  $\mathcal{M}$  in a fts  $(X, \tau)$  such that  $x_\lambda q\mathcal{M} \leq \mathcal{A}$  (resp.  $\mathcal{B}q\mathcal{M} \leq \mathcal{A}$ ).

**Theorem 3.15:** A fuzzy set  $\mathcal{A}$  of a fts  $(X, \tau)$  is  $F\alpha^m$ -CS iff  $\mathcal{A}\bar{q}\mathcal{K} \Rightarrow \text{int}(\text{cl}(\mathcal{A}))\bar{q}\mathcal{K}$ , for every  $F\alpha$ -CS  $\mathcal{K}$  of  $X$ .

**Proof: Necessity.** Let  $\mathcal{K}$  be a  $F\alpha$ -CS and  $\mathcal{A}\bar{q}\mathcal{K}$ . Then  $\mathcal{A} \leq 1_X - \mathcal{K}$  and  $1_X - \mathcal{K}$  is a  $F\alpha$ -OS in  $X$  which implies that  $\text{int}(\text{cl}(\mathcal{A})) \leq 1_X - \mathcal{K}$  as  $\mathcal{A}$  is a  $F\alpha^m$ -CS. Hence,  $\text{int}(\text{cl}(\mathcal{A}))\bar{q}\mathcal{K}$ .

**Sufficiency.** Let  $\mathcal{U}$  be a  $F\alpha$ -OS of a fts  $(X, \tau)$  such that  $\mathcal{A} \leq \mathcal{U}$ . Then  $\mathcal{A}\bar{q}(1_X - \mathcal{U})$  and  $1_X - \mathcal{U}$  is a  $F\alpha$ -CS in  $X$ . By hypothesis,  $\text{int}(\text{cl}(\mathcal{A}))\bar{q}(1_X - \mathcal{U})$  implies  $\text{int}(\text{cl}(\mathcal{A})) \leq \mathcal{U}$ . Hence,  $\mathcal{A}$  is a  $F\alpha^m$ -CS in  $X$ .

**Theorem 3.16:** Let  $x_\lambda$  and  $\mathcal{A}$  be a fuzzy point and a fuzzy set respectively in a fts  $(X, \tau)$ . Then  $x_\lambda \in \alpha^m\text{-cl}(\mathcal{A})$  iff every  $F\alpha^m$ - $q$ -nhd of  $x_\lambda$  is  $q$ -coincident with  $\mathcal{A}$ .

**Proof:** Let  $x_\lambda \in \alpha^m\text{-cl}(\mathcal{A})$ . Suppose there exists a  $F\alpha^m$ - $q$ -nhd  $\mathcal{M}$  of  $x_\lambda$  such that  $\mathcal{M}\bar{q}\mathcal{A}$ . Since  $\mathcal{M}$  is a  $F\alpha^m$ - $q$ -nhd of  $x_\lambda$ , there exists a  $F\alpha^m$ -OS  $\mathcal{N}$  in  $X$  such that  $x_\lambda q\mathcal{N} \leq \mathcal{M}$  which gives that  $\mathcal{N}\bar{q}\mathcal{A}$  and hence  $\mathcal{A} \leq 1_X - \mathcal{N}$ . Then  $\alpha^m\text{-cl}(\mathcal{A}) \leq 1_X - \mathcal{N}$ , as  $1_X - \mathcal{N}$  is a  $F\alpha^m$ -CS. Since  $x_\lambda \notin 1_X - \mathcal{N}$ , we have  $x_\lambda \notin \alpha^m\text{-cl}(\mathcal{A})$ , a contradiction. Thus every  $F\alpha^m$ - $q$ -nhd of  $x_\lambda$  is  $q$ -coincident with  $\mathcal{A}$ .

Conversely, suppose  $x_\lambda \notin \alpha^m\text{-cl}(\mathcal{A})$ . Then there exists a  $F\alpha^m$ -CS  $\mathcal{B}$  such that  $\mathcal{A} \leq \mathcal{B}$  and  $x_\lambda \notin \mathcal{B}$ . Then we have  $x_\lambda q(1_X - \mathcal{B})$  and  $\mathcal{A}\bar{q}(1_X - \mathcal{B})$ , a contradiction. Hence  $x_\lambda \in \alpha^m\text{-cl}(\mathcal{A})$ .

**Theorem 3.17:** Let  $\mathcal{A}$  and  $\mathcal{B}$  be two fuzzy sets in a fts  $(X, \tau)$ . Then the following are true:

$$(i) \alpha^m\text{-cl}(0_X) = 0_X, \alpha^m\text{-cl}(1_X) = 1_X.$$

$$(ii) \alpha^m\text{-cl}(\mathcal{A}) \text{ is a } F\alpha^m\text{-CS in } X.$$

$$(iii) \alpha^m\text{-cl}(\mathcal{A}) \leq \alpha^m\text{-cl}(\mathcal{B}) \text{ when } \mathcal{A} \leq \mathcal{B}.$$

$$(iv) \mathcal{M}q\mathcal{A} \text{ iff } \mathcal{M}q\alpha^m\text{-cl}(\mathcal{A}), \text{ when } \mathcal{M} \text{ is a } F\alpha^m\text{-OS in } X.$$

$$(v) \alpha^m\text{-cl}(\mathcal{A}) = \alpha^m\text{-cl}(\alpha^m\text{-cl}(\mathcal{A})).$$

**Proof:** (i) and (ii) are obvious.

(iii) Let  $x_\lambda \notin \alpha^m\text{-cl}(\mathcal{B})$ . By theorem (3.16), there is a  $F\alpha^m$ - $q$ -nhd  $\mathcal{N}$  of a fuzzy point  $x_\lambda$  such that  $\mathcal{N}\bar{q}\mathcal{B}$ , so there is a  $F\alpha^m$ -OS  $\mathcal{M}$  such that  $x_\lambda q\mathcal{M} \leq \mathcal{N}$  and  $\mathcal{M}\bar{q}\mathcal{B}$ . Since  $\mathcal{A} \leq \mathcal{B}$ , then  $\mathcal{M}\bar{q}\mathcal{A}$ . Hence  $x_\lambda \notin \alpha^m\text{-cl}(\mathcal{A})$  by theorem (3.16). Thus  $\alpha^m\text{-cl}(\mathcal{A}) \leq \alpha^m\text{-cl}(\mathcal{B})$ .

(iv) Let  $\mathcal{M}$  be a  $F\alpha^m$ -OS in  $X$ . Suppose that  $\mathcal{M}\bar{q}\mathcal{A}$ , then  $\mathcal{A} \leq 1_X - \mathcal{M}$ . Since  $1_X - \mathcal{M}$  is a  $F\alpha^m$ -CS and by a part (iii),  $\alpha^m\text{-cl}(\mathcal{A}) \leq \alpha^m\text{-cl}(1_X - \mathcal{M}) = 1_X - \mathcal{M}$ . Hence,  $\mathcal{M}\bar{q}\alpha^m\text{-cl}(\mathcal{A})$ .

Conversely, suppose that  $\mathcal{M}\bar{q}\alpha^m\text{-cl}(\mathcal{A})$ . Then  $\alpha^m\text{-cl}(\mathcal{A}) \leq 1_X - \mathcal{M}$ . Since  $\mathcal{A} \leq \alpha^m\text{-cl}(\mathcal{A})$ , we have  $\mathcal{A} \leq 1_X - \mathcal{M}$ . Hence  $\mathcal{M}\bar{q}\mathcal{A}$ .


(v) Since  $\alpha^m\text{-cl}(\mathcal{A}) \leq \alpha^m\text{-cl}(\alpha^m\text{-cl}(\mathcal{A}))$ . We prove that  $\alpha^m\text{-cl}(\alpha^m\text{-cl}(\mathcal{A})) \leq \alpha^m\text{-cl}(\mathcal{A})$ . Suppose that  $x_\lambda \notin \alpha^m\text{-cl}(\mathcal{A})$ . Then by theorem (3.16), there exists a  $F\alpha^m$ -q-nhd  $\mathcal{N}$  of a fuzzy point  $x_\lambda$  such that  $\mathcal{N} \bar{q} \mathcal{A}$  and so there is a  $F\alpha^m$ -OS  $\mathcal{M}$  in  $X$  such that  $x_\lambda q \mathcal{M} \leq \mathcal{N}$  and  $\mathcal{M} \bar{q} \mathcal{A}$ . By a part (iv),  $\mathcal{M} \bar{q} \alpha^m\text{-cl}(\mathcal{A})$ . Then by theorem (3.16),  $x_\lambda \notin \alpha^m\text{-cl}(\alpha^m\text{-cl}(\mathcal{A}))$ . Thus  $\alpha^m\text{-cl}(\alpha^m\text{-cl}(\mathcal{A})) \leq \alpha^m\text{-cl}(\mathcal{A})$ . Hence  $\alpha^m\text{-cl}(\mathcal{A}) = \alpha^m\text{-cl}(\alpha^m\text{-cl}(\mathcal{A}))$ .

**Theorem 3.18:** Let  $\mathcal{A}$  and  $\mathcal{B}$  be two fuzzy sets in a fts  $(X, \tau)$ . Then the following are true:

- (i)  $\alpha^m\text{-int}(0_X) = 0_X$ ,  $\alpha^m\text{-int}(1_X) = 1_X$ .
- (ii)  $\alpha^m\text{-int}(\mathcal{A})$  is a  $F\alpha^m$ -OS in  $X$ .
- (iii)  $\alpha^m\text{-int}(\mathcal{A}) \leq \alpha^m\text{-int}(\mathcal{B})$  when  $\mathcal{A} \leq \mathcal{B}$ .
- (iv)  $\alpha^m\text{-int}(\mathcal{A}) = \alpha^m\text{-int}(\alpha^m\text{-int}(\mathcal{A}))$ .

**Proof:** Obvious.

**Remark 3.19:** The following are the implications of a  $F\alpha^m$ -CS and the reverse is not true.



#### 4. FUZZY $\alpha^m$ -KERNEL AND FUZZY $\alpha^m$ - $R_i$ -SPACES, $i = 0, 1, 2, 3$

**Definition 4.1:** The intersection of all  $F\alpha^m$ -open subset of  $X$  containing  $\mathcal{M}$  is called the fuzzy  $\alpha^m$ -kernel of  $\mathcal{M}$  (briefly  $\alpha^m\text{-ker}(\mathcal{M})$ ), this means  $\alpha^m\text{-ker}(\mathcal{M}) = \bigwedge \{ \mathcal{U} \in F\alpha^m\text{-O}(X) : \mathcal{M} \leq \mathcal{U} \}$ .

**Definition 4.2:** In a fts  $(X, \tau)$ , a fuzzy set  $\mathcal{M}$  is said to be weakly ultra fuzzy  $\alpha^m$ -separated from  $\mathcal{N}$  if there exists a  $F\alpha^m$ -OS  $\mathcal{U}$  such that  $\mathcal{U} \wedge \mathcal{N} = 0_X$  or  $\mathcal{M} \wedge \alpha^m\text{-cl}(\mathcal{N}) = 0_X$ .

By definition (4.2), we have the following: For every two distinct fuzzy points  $x_\lambda$  and  $y_\sigma$  of  $X$ ,

- (i)  $\alpha^m\text{-cl}(\{x_\lambda\}) = \{y_\sigma : \{y_\sigma\} \text{ is not weakly ultra fuzzy } \alpha^m\text{-separated from } \{x_\lambda\}\}$ .
- (ii)  $\alpha^m\text{-ker}(\{x_\lambda\}) = \{y_\sigma : \{x_\lambda\} \text{ is not weakly ultra fuzzy } \alpha^m\text{-separated from } \{y_\sigma\}\}$ .

**Corollary 4.3:** Let  $(X, \tau)$  be a fts, then  $y_\sigma \in \alpha^m\text{-ker}(\{x_\lambda\})$  iff  $x_\lambda \in \alpha^m\text{-cl}(\{y_\sigma\})$  for each  $x \neq y \in X$ .

**Proof:** Suppose that  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$ . Then there exists a  $F\alpha^m$ -OS  $\mathcal{U}$  containing  $x_\lambda$  such that  $y_\sigma \notin \mathcal{U}$ . Therefore, we have  $x_\lambda \notin \alpha^m\text{-cl}(\{y_\sigma\})$ . The converse part can be proved in a similar way.

**Definition 4.4:** A fts  $(X, \tau)$  is called fuzzy  $\alpha^m$ - $R_0$ -space ( $F\alpha^m$ - $R_0$ -space, for short) if for each  $F\alpha^m$ -OS  $\mathcal{U}$  and  $x_\lambda \in \mathcal{U}$ , then  $\alpha^m\text{-cl}(\{x_\lambda\}) \leq \mathcal{U}$ .

**Definition 4.5:** A fts  $(X, \tau)$  is called fuzzy  $\alpha^m$ - $R_1$ -space ( $F\alpha^m$ - $R_1$ -space, for short) if for each two distinct fuzzy points  $x_\lambda$  and  $y_\sigma$  of  $X$  with  $\alpha^m\text{-cl}(\{x_\lambda\}) \neq \alpha^m\text{-cl}(\{y_\sigma\})$ , there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\alpha^m\text{-cl}(\{x_\lambda\}) \leq \mathcal{U}$  and  $\alpha^m\text{-cl}(\{y_\sigma\}) \leq \mathcal{V}$ .

**Theorem 4.6:** Let  $(X, \tau)$  be a fts. Then  $(X, \tau)$  is  $F\alpha^m$ - $R_0$ -space iff  $\alpha^m\text{-cl}(\{x_\lambda\}) = \alpha^m\text{-ker}(\{x_\lambda\})$ , for each  $x \in X$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-R_0$ -space. If  $\alpha^m-cl(\{x_\lambda\}) \neq \alpha^m-ker(\{x_\lambda\})$ , for each  $x \in X$ , then there exist another fuzzy point  $y \neq x$  such that  $y_\sigma \in \alpha^m-cl(\{x_\lambda\})$  and  $y_\sigma \notin \alpha^m-ker(\{x_\lambda\})$  this means there exist an  $\mathcal{U}_{x_\lambda}$   $F\alpha^m$ -OS,  $y_\sigma \notin \mathcal{U}_{x_\lambda}$  implies  $\alpha^m-cl(\{x_\lambda\}) \not\leq \mathcal{U}_{x_\lambda}$  this contradiction. Thus  $\alpha^m-cl(\{x_\lambda\}) = \alpha^m-ker(\{x_\lambda\})$ .

Conversely, let  $\alpha^m-cl(\{x_\lambda\}) = \alpha^m-ker(\{x_\lambda\})$ , for each  $F\alpha^m$ -OS  $\mathcal{U}, x_\lambda \in \mathcal{U}$ , then  $\alpha^m-ker(\{x_\lambda\}) = \alpha^m-cl(\{x_\lambda\}) \leq \mathcal{U}$  [by definition (4.1)]. Hence by definition (4.4),  $(X, \tau)$  is a  $F\alpha^m-R_0$ -space.

**Theorem 4.7:** A fts  $(X, \tau)$  is an  $F\alpha^m-R_0$ -space iff for each  $\mathcal{A}$   $F\alpha^m$ -CS and  $x_\lambda \in \mathcal{A}$ , then  $\alpha^m-ker(\{x_\lambda\}) \leq \mathcal{A}$ .

**Proof:** Let for each  $\mathcal{A}$   $F\alpha^m$ -CS and  $x_\lambda \in \mathcal{A}$ , then  $\alpha^m-ker(\{x_\lambda\}) \leq \mathcal{A}$  and let  $\mathcal{U}$  be a  $F\alpha^m$ -OS,  $x_\lambda \in \mathcal{U}$  then for each  $y_\sigma \notin \mathcal{U}$  implies  $y_\sigma \in \mathcal{U}^c$  is a  $F\alpha^m$ -CS implies  $\alpha^m-ker(\{y_\sigma\}) \leq \mathcal{U}^c$  [by assumption]. Therefore  $x_\lambda \notin \alpha^m-ker(\{y_\sigma\})$  implies  $y_\sigma \notin \alpha^m-cl(\{x_\lambda\})$  [by corollary (4.3)]. So  $\alpha^m-cl(\{x_\lambda\}) \leq \mathcal{U}$ . Thus  $(X, \tau)$  is an  $F\alpha^m-R_0$ -space.

Conversely, let  $(X, \tau)$  be a  $F\alpha^m-R_0$ -space and  $\mathcal{A}$  be a  $F\alpha^m$ -CS and  $x_\lambda \in \mathcal{A}$ . Then for each  $y_\sigma \notin \mathcal{A}$  implies  $y_\sigma \in \mathcal{A}^c$  is a  $F\alpha^m$ -OS, then  $\alpha^m-cl(\{y_\sigma\}) \leq \mathcal{A}^c$  [since  $(X, \tau)$  is a  $F\alpha^m-R_0$ -space], so  $\alpha^m-ker(\{x_\lambda\}) = \alpha^m-cl(\{x_\lambda\})$ . Thus  $\alpha^m-ker(\{x_\lambda\}) \leq \mathcal{A}$ .

**Corollary 4.8:** A fts  $(X, \tau)$  is  $F\alpha^m-R_0$ -space iff for each  $\mathcal{U}$   $F\alpha^m$ -OS and  $x_\lambda \in \mathcal{U}$ , then  $\alpha^m-cl(\alpha^m-ker(\{x_\lambda\})) \leq \mathcal{U}$ .

**Proof:** Clearly.

**Theorem 4.9:** Every  $F\alpha^m-R_1$ -space is a  $F\alpha^m-R_0$ -space.

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-R_1$ -space and let  $\mathcal{U}$  be a  $F\alpha^m$ -OS,  $x_\lambda \in \mathcal{U}$ , then for each  $y_\sigma \notin \mathcal{U}$  implies  $y_\sigma \in \mathcal{U}^c$  is a  $F\alpha^m$ -CS and  $\alpha^m-cl(\{y_\sigma\}) \leq \mathcal{U}^c$  implies  $\alpha^m-cl(\{x_\lambda\}) \neq \alpha^m-cl(\{y_\sigma\})$ . Hence by definition (4.5),  $\alpha^m-cl(\{x_\lambda\}) \leq \mathcal{U}$ . Thus  $(X, \tau)$  is a  $F\alpha^m-R_0$ -space.

**Theorem 4.10:** A fts  $(X, \tau)$  is  $F\alpha^m-R_1$ -space iff for each  $x \neq y \in X$  with  $\alpha^m-ker(\{x_\lambda\}) \neq \alpha^m-ker(\{y_\sigma\})$ , then there exist  $F\alpha^m$ -CS  $\mathcal{A}_1, \mathcal{A}_2$  such that  $\alpha^m-ker(\{x_\lambda\}) \leq \mathcal{A}_1$ ,  $\alpha^m-ker(\{x_\lambda\}) \wedge \mathcal{A}_2 = 0_X$  and  $\alpha^m-ker(\{y_\sigma\}) \leq \mathcal{A}_2$ ,  $\alpha^m-ker(\{y_\sigma\}) \wedge \mathcal{A}_1 = 0_X$  and  $\mathcal{A}_1 \vee \mathcal{A}_2 = 1_X$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-R_1$ -space. Then for each  $x \neq y \in X$  with  $\alpha^m-ker(\{x_\lambda\}) \neq \alpha^m-ker(\{y_\sigma\})$ . Since every  $F\alpha^m-R_1$ -space is a  $F\alpha^m-R_0$ -space [by theorem (4.9)], and by theorem (4.6),  $\alpha^m-cl(\{x_\lambda\}) \neq \alpha^m-cl(\{y_\sigma\})$ , then there exist  $F\alpha^m$ -OS  $\mathcal{U}_1, \mathcal{U}_2$  such that  $\alpha^m-cl(\{x_\lambda\}) \leq \mathcal{U}_1$  and  $\alpha^m-cl(\{y_\sigma\}) \leq \mathcal{U}_2$  and  $\mathcal{U}_1 \wedge \mathcal{U}_2 = 0_X$  [since  $(X, \tau)$  is a  $F\alpha^m-R_1$ -space], then  $\mathcal{U}_1^c$  and  $\mathcal{U}_2^c$  are  $F\alpha^m$ -CS such that  $\mathcal{U}_1^c \vee \mathcal{U}_2^c = 1_X$ . Put  $\mathcal{A}_1 = \mathcal{U}_1^c$  and  $\mathcal{A}_2 = \mathcal{U}_2^c$ . Thus  $x_\lambda \in \mathcal{U}_1 \leq \mathcal{A}_2$  and  $y_\sigma \in \mathcal{U}_2 \leq \mathcal{A}_1$  so that  $\alpha^m-ker(\{x_\lambda\}) \leq \mathcal{U}_1 \leq \mathcal{A}_2$  and  $\alpha^m-ker(\{y_\sigma\}) \leq \mathcal{U}_2 \leq \mathcal{A}_1$ .

Conversely, let for each  $x \neq y \in X$  with  $\alpha^m-ker(\{x_\lambda\}) \neq \alpha^m-ker(\{y_\sigma\})$ , there exist  $F\alpha^m$ -CS  $\mathcal{A}_1, \mathcal{A}_2$  such that  $\alpha^m-ker(\{x_\lambda\}) \leq \mathcal{A}_1$ ,  $\alpha^m-ker(\{x_\lambda\}) \wedge \mathcal{A}_2 = 0_X$  and  $\alpha^m-ker(\{y_\sigma\}) \leq \mathcal{A}_2$ ,  $\alpha^m-ker(\{y_\sigma\}) \wedge \mathcal{A}_1 = 0_X$  and  $\mathcal{A}_1 \vee \mathcal{A}_2 = 1_X$ , then  $\mathcal{A}_1^c$  and  $\mathcal{A}_2^c$  are  $F\alpha^m$ -OS such that  $\mathcal{A}_1^c \wedge \mathcal{A}_2^c = 0_X$ . Put  $\mathcal{U}_1 = \mathcal{A}_1^c$  and  $\mathcal{U}_2 = \mathcal{A}_2^c$ . Thus,  $\alpha^m-ker(\{x_\lambda\}) \leq \mathcal{U}_1$  and  $\alpha^m-ker(\{y_\sigma\}) \leq \mathcal{U}_2$  and  $\mathcal{U}_1 \wedge \mathcal{U}_2 = 0_X$ , so that  $x_\lambda \in \mathcal{U}_1$  and  $y_\sigma \in \mathcal{U}_2$  implies  $x_\lambda \notin \alpha^m-cl(\{y_\sigma\})$  and  $y_\sigma \notin \alpha^m-cl(\{x_\lambda\})$ , then  $\alpha^m-cl(\{x_\lambda\}) \leq \mathcal{U}_1$  and  $\alpha^m-cl(\{y_\sigma\}) \leq \mathcal{U}_2$ . Thus,  $(X, \tau)$  is a  $F\alpha^m-R_1$ -space.

**Corollary 4.11:** A fts  $(X, \tau)$  is  $F\alpha^m-R_1$ -space iff for each  $x \neq y \in X$  with  $\alpha^m-cl(\{x_\lambda\}) \neq \alpha^m-cl(\{y_\sigma\})$  there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\alpha^m-cl(\alpha^m-ker(\{x_\lambda\})) \leq \mathcal{U}$  and  $\alpha^m-cl(\alpha^m-ker(\{y_\sigma\})) \leq \mathcal{V}$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-R_1$ -space and let  $x \neq y \in X$  with  $\alpha^m-cl(\{x_\lambda\}) \neq \alpha^m-cl(\{y_\sigma\})$ , then there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\alpha^m-cl(\{x_\lambda\}) \leq \mathcal{U}$  and  $\alpha^m-cl(\{y_\sigma\}) \leq \mathcal{V}$ . Also  $(X, \tau)$  is  $F\alpha^m-R_0$ -space [by theorem (4.9)] implies for each  $x \in X$ , then  $\alpha^m-cl(\{x_\lambda\}) = \alpha^m-ker(\{x_\lambda\})$  [by theorem (4.6)], but  $\alpha^m-cl(\{x_\lambda\}) = \alpha^m-cl(\alpha^m-cl(\{x_\lambda\})) = \alpha^m-cl(\alpha^m-ker(\{x_\lambda\}))$ . Thus  $\alpha^m-cl(\alpha^m-ker(\{x_\lambda\})) \leq \mathcal{U}$  and  $\alpha^m-cl(\alpha^m-ker(\{y_\sigma\})) \leq \mathcal{V}$ .

Conversely, let for each  $x \neq y \in X$  with  $\alpha^m-cl(\{x_\lambda\}) \neq \alpha^m-cl(\{y_\sigma\})$  there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\alpha^m-cl(\alpha^m-ker(\{x_\lambda\})) \leq \mathcal{U}$  and  $\alpha^m-cl(\alpha^m-ker(\{y_\sigma\})) \leq \mathcal{V}$ . Since  $\{x_\lambda\} \leq \alpha^m-ker(\{x_\lambda\})$ , then  $\alpha^m-cl(\{x_\lambda\}) \leq \alpha^m-cl(\alpha^m-ker(\{x_\lambda\}))$  for each  $x \in X$ . So we get  $\alpha^m-cl(\{x_\lambda\}) \leq \mathcal{U}$  and  $\alpha^m-cl(\{y_\sigma\}) \leq \mathcal{V}$ . Thus,  $(X, \tau)$  is a  $F\alpha^m-R_1$ -space.



**Definition 4.12:** Let  $(X, \tau)$  be a fts. Then  $X$  is called:

- (i) fuzzy  $\alpha^m$ -regular space ( $F\alpha^m r$ -space, for short), if for each fuzzy point  $x_\lambda$  and each  $F\alpha^m$ -CS  $\mathcal{F}$  such that  $x_\lambda \in 1_X - \mathcal{F}$ , there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}$  and  $\mathcal{V}$  such that  $x_\lambda \in \mathcal{U}$  and  $\mathcal{F} \leq \mathcal{V}$ .
- (ii) fuzzy  $\alpha^m$ -normal space ( $F\alpha^m n$ -space, for short) iff for each pair of disjoint  $F\alpha^m$ -CS  $\mathcal{A}$  and  $\mathcal{B}$ , there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}$  and  $\mathcal{V}$  such that  $\mathcal{A} \leq \mathcal{U}$  and  $\mathcal{B} \leq \mathcal{V}$ .
- (iii) fuzzy  $\alpha^m$ - $R_2$ -space ( $F\alpha^m R_2$ -space, for short) if it is property  $F\alpha^m r$ -space.
- (iv) fuzzy  $\alpha^m$ - $R_3$ -space ( $F\alpha^m R_3$ -space, for short) iff it is  $F\alpha^m R_1$ -space and  $F\alpha^m n$ -space.

**Example 4.13:** Consider the fts  $(X, \tau)$  of example (3.2). Then  $(X, \tau)$  is a  $F\alpha^m r$ -space and  $F\alpha^m n$ -space.

**Remark 4.14:** Every  $F\alpha^m R_k$ -space is a  $F\alpha^m R_{k-1}$ -space,  $k = 2, 3$ .

**Proof:** Clearly.

**Theorem 4.15:** A fts  $(X, \tau)$  is  $F\alpha^m r$ -space ( $F\alpha^m R_2$ -space) iff for each  $F\alpha^m$ -closed subset  $\mathcal{A}$  of  $X$  and  $x_\lambda \notin \mathcal{A}$  with  $\alpha^m\text{-ker}(\mathcal{A}) \neq \alpha^m\text{-ker}(\{x_\lambda\})$  then there exist  $F\alpha^m$ -CS  $\mathcal{F}_1, \mathcal{F}_2$  such that  $\alpha^m\text{-ker}(\mathcal{A}) \leq \mathcal{F}_1$ ,  $\alpha^m\text{-ker}(\mathcal{A}) \wedge \mathcal{F}_2 = 0_X$  and  $\alpha^m\text{-ker}(\{x_\lambda\}) \leq \mathcal{F}_2$ ,  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \mathcal{F}_1 = 0_X$  and  $\mathcal{F}_1 \vee \mathcal{F}_2 = 1_X$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m r$ -space ( $F\alpha^m R_2$ -space) and let  $\mathcal{A}$  be a  $F\alpha^m$ -CS,  $x_\lambda \notin \mathcal{A}$ , then there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\mathcal{A} \leq \mathcal{U}$ ,  $x_\lambda \in \mathcal{V}$  and  $\mathcal{U} \wedge \mathcal{V} = 0_X$ , then  $\mathcal{U}^c$  and  $\mathcal{V}^c$  are  $F\alpha^m$ -CS such that  $\mathcal{U}^c \vee \mathcal{V}^c = 1_X$ .

Put  $\mathcal{F}_2 = \mathcal{U}^c$  and  $\mathcal{F}_1 = \mathcal{V}^c$ , so we get  $\alpha^m\text{-ker}(\mathcal{A}) \leq \mathcal{U} \leq \mathcal{F}_1$ ,  $\alpha^m\text{-ker}(\mathcal{A}) \wedge \mathcal{F}_2 = 0_X$  and  $\alpha^m\text{-ker}(\{x_\lambda\}) \leq \mathcal{V} \leq \mathcal{F}_2$ ,  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \mathcal{F}_1 = 0_X$  and  $\mathcal{F}_1 \vee \mathcal{F}_2 = 1_X$ .

Conversely, let for each  $F\alpha^m$ -closed subset  $\mathcal{A}$  of  $X$  and  $x_\lambda \notin \mathcal{A}$  with  $\alpha^m\text{-ker}(\mathcal{A}) \neq \alpha^m\text{-ker}(\{x_\lambda\})$ , then there exist  $F\alpha^m$ -CS  $\mathcal{F}_1, \mathcal{F}_2$  such that  $\alpha^m\text{-ker}(\mathcal{A}) \leq \mathcal{F}_1$ ,  $\alpha^m\text{-ker}(\mathcal{A}) \wedge \mathcal{F}_2 = 0_X$  and  $\alpha^m\text{-ker}(\{x_\lambda\}) \leq \mathcal{F}_2$ ,  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \mathcal{F}_1 = 0_X$  and  $\mathcal{F}_1 \vee \mathcal{F}_2 = 1_X$ . Then  $\mathcal{F}_1^c$  and  $\mathcal{F}_2^c$  are  $F\alpha^m$ -OS such that  $\mathcal{F}_1^c \wedge \mathcal{F}_2^c = 0_X$  and  $\alpha^m\text{-ker}(\mathcal{A}) \wedge \mathcal{F}_1^c = 0_X$ ,  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \mathcal{F}_2^c = 0_X$ . So that  $\mathcal{A} \leq \mathcal{F}_2^c$  and  $x_\lambda \in \mathcal{F}_1^c$ . Thus,  $(X, \tau)$  is a  $F\alpha^m r$ -space ( $F\alpha^m R_2$ -space).

**Lemma 4.16:** Let  $(X, \tau)$  be a  $F\alpha^m r$ -space and  $\mathcal{F}$  be a  $F\alpha^m$ -CS. Then  $\alpha^m\text{-ker}(\mathcal{F}) = \mathcal{F} = \alpha^m\text{-cl}(\mathcal{F})$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m r$ -space and  $\mathcal{F}$  be a  $F\alpha^m$ -CS. Then for each  $x_\lambda \notin \mathcal{F}$ , there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\mathcal{F} \leq \mathcal{U}$  and  $x_\lambda \in \mathcal{V}$ . Since  $\alpha^m\text{-ker}(\mathcal{F}) \leq \mathcal{U}$ , implies  $\alpha^m\text{-ker}(\mathcal{F}) \wedge \mathcal{V} = 0_X$ , thus  $x_\lambda \notin \alpha^m\text{-cl}(\alpha^m\text{-ker}(\mathcal{F}))$ . We showing that if  $x_\lambda \notin \mathcal{F}$  implies  $x_\lambda \notin \alpha^m\text{-cl}(\alpha^m\text{-ker}(\mathcal{F}))$ , therefore  $\alpha^m\text{-cl}(\alpha^m\text{-ker}(\mathcal{F})) \leq \mathcal{F} = \alpha^m\text{-cl}(\mathcal{F})$ . As  $\alpha^m\text{-cl}(\mathcal{F}) = \mathcal{F} \leq \alpha^m\text{-ker}(\mathcal{F})$  [by definition (4.1)]. Thus,  $\alpha^m\text{-ker}(\mathcal{F}) = \mathcal{F} = \alpha^m\text{-cl}(\mathcal{F})$ .

**Theorem 4.17:** A fts  $(X, \tau)$  is  $F\alpha^m r$ -space ( $F\alpha^m R_2$ -space) iff for each  $F\alpha^m$ -closed subset  $\mathcal{F}$  of  $X$  and  $x_\lambda \notin \mathcal{F}$  with  $\alpha^m\text{-cl}(\alpha^m\text{-ker}(\mathcal{F})) \neq \alpha^m\text{-cl}(\alpha^m\text{-ker}(\{x_\lambda\}))$ , then there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\alpha^m\text{-cl}(\alpha^m\text{-ker}(\mathcal{F})) \leq \mathcal{U}$  and  $\alpha^m\text{-cl}(\alpha^m\text{-ker}(\{x_\lambda\})) \leq \mathcal{V}$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m r$ -space ( $F\alpha^m R_2$ -space) and let  $\mathcal{F}$  be a  $F\alpha^m$ -CS,  $x_\lambda \notin \mathcal{F}$ . Then there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\mathcal{F} \leq \mathcal{U}$  and  $x_\lambda \in \mathcal{V}$ . By lemma (4.16),  $\alpha^m\text{-cl}(\alpha^m\text{-ker}(\mathcal{F})) = \alpha^m\text{-cl}(\mathcal{F}) = \mathcal{F}$ , in the other hand  $(X, \tau)$  is a  $F\alpha^m R_0$ -space [by theorem (4.9) and remark (4.14)]. Hence, by theorem (4.6),  $\alpha^m\text{-cl}(\{x_\lambda\}) = \alpha^m\text{-ker}(\{x_\lambda\})$ , for each  $x \in X$ . Thus,  $\alpha^m\text{-cl}(\alpha^m\text{-ker}(\mathcal{F})) \leq \mathcal{U}$  and  $\alpha^m\text{-cl}(\alpha^m\text{-ker}(\{x_\lambda\})) \leq \mathcal{V}$ .

Conversely, let for each  $F\alpha^m$ -CS  $\mathcal{F}$  and  $x_\lambda \notin \mathcal{F}$  with  $\alpha^m\text{-cl}(\alpha^m\text{-ker}(\mathcal{F})) \neq \alpha^m\text{-cl}(\alpha^m\text{-ker}(\{x_\lambda\}))$ , then there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\alpha^m\text{-cl}(\alpha^m\text{-ker}(\mathcal{F})) \leq \mathcal{U}$  and  $\alpha^m\text{-cl}(\alpha^m\text{-ker}(\{x_\lambda\})) \leq \mathcal{V}$ . Then  $\mathcal{F} \leq \mathcal{U}$  and  $x_\lambda \in \mathcal{V}$ . Thus,  $(X, \tau)$  is a  $F\alpha^m r$ -space ( $F\alpha^m R_2$ -space).

**Theorem 4.18:** A fts  $(X, \tau)$  is  $F\alpha^m n$ -space iff for each disjoint  $F\alpha^m$ -CS  $\mathcal{A}, \mathcal{B}$  with  $\alpha^m\text{-ker}(\mathcal{A}) \neq \alpha^m\text{-ker}(\mathcal{B})$  then there exist  $F\alpha^m$ -CS  $\mathcal{F}_1, \mathcal{F}_2$  such that  $\alpha^m\text{-ker}(\mathcal{A}) \leq \mathcal{F}_1$ ,  $\alpha^m\text{-ker}(\mathcal{A}) \wedge \mathcal{F}_2 = 0_X$  and  $\alpha^m\text{-ker}(\mathcal{B}) \leq \mathcal{F}_2$ ,  $\alpha^m\text{-ker}(\mathcal{B}) \wedge \mathcal{F}_1 = 0_X$  and  $\mathcal{F}_1 \vee \mathcal{F}_2 = 1_X$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m$ - $n$ -space and let for each disjoint  $F\alpha^m$ -CS  $\mathcal{A}, \mathcal{B}$  with  $\alpha^m\text{-ker}(\mathcal{A}) \neq \alpha^m\text{-ker}(\mathcal{B})$  then there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\mathcal{A} \leq \mathcal{U}$  and  $\mathcal{B} \leq \mathcal{V}$  and  $\mathcal{U} \wedge \mathcal{V} = 0_X$ , then  $\mathcal{U}^c$  and  $\mathcal{V}^c$  are  $F\alpha^m$ -CS such that  $\mathcal{U}^c \vee \mathcal{V}^c = 1_X$  and  $\alpha^m\text{-ker}(\mathcal{A}) \wedge \mathcal{U}^c = 0_X$ ,  $\alpha^m\text{-ker}(\mathcal{B}) \wedge \mathcal{V}^c = 0_X$  Put  $\mathcal{U}^c = \mathcal{F}_2$  and  $\mathcal{V}^c = \mathcal{F}_1$ . Thus,  $\alpha^m\text{-ker}(\mathcal{A}) \leq \mathcal{F}_1$ ,  $\alpha^m\text{-ker}(\mathcal{A}) \wedge \mathcal{F}_2 = 0_X$  and  $\alpha^m\text{-ker}(\mathcal{B}) \leq \mathcal{F}_2$ ,  $\alpha^m\text{-ker}(\mathcal{B}) \wedge \mathcal{F}_1 = 0_X$ .

Conversely, let for each disjoint  $F\alpha^m$ -CS  $\mathcal{A}, \mathcal{B}$  with  $\alpha^m\text{-ker}(\mathcal{A}) \neq \alpha^m\text{-ker}(\mathcal{B})$ , there exist  $F\alpha^m$ -CS  $\mathcal{F}_1, \mathcal{F}_2$  such that  $\alpha^m\text{-ker}(\mathcal{A}) \leq \mathcal{F}_1$ ,  $\alpha^m\text{-ker}(\mathcal{A}) \wedge \mathcal{F}_2 = 0_X$  and  $\alpha^m\text{-ker}(\mathcal{B}) \leq \mathcal{F}_2$ ,  $\alpha^m\text{-ker}(\mathcal{B}) \wedge \mathcal{F}_1 = 0_X$  and  $\mathcal{F}_1 \vee \mathcal{F}_2 = 1_X$  implies  $\mathcal{F}_1^c$  and  $\mathcal{F}_2^c$  are  $F\alpha^m$ -OS such that  $\mathcal{F}_1^c \wedge \mathcal{F}_2^c = 0_X$ . Put  $\mathcal{F}_1^c = \mathcal{V}$  and  $\mathcal{F}_2^c = \mathcal{U}$ , thus  $\alpha^m\text{-ker}(\mathcal{A}) \leq \mathcal{U}$  and  $\alpha^m\text{-ker}(\mathcal{B}) \leq \mathcal{V}$ , so that  $\mathcal{A} \leq \mathcal{U}$  and  $\mathcal{B} \leq \mathcal{V}$ . Thus  $(X, \tau)$  is a  $F\alpha^m$ - $n$ -space.

**Theorem 4.19:** Every  $F\alpha^m$ - $R_3$ -space is a  $F\alpha^m$ - $r$ -space.

**Proof:** Let  $\mathcal{F}$  be a  $F\alpha^m$ -CS and  $x_\lambda \notin \mathcal{F}$ . Then  $\alpha^m\text{-ker}(\{x_\lambda\}) \neq \alpha^m\text{-ker}(\mathcal{F})$ , then for each  $y_\sigma \in \mathcal{F}$  there exist  $F\alpha^m$ -CS  $\mathcal{A}_{y_\sigma}, \mathcal{B}_{y_\sigma}$  such that  $\alpha^m\text{-ker}(\{y_\sigma\}) \leq \mathcal{A}_{y_\sigma}$ ,  $\alpha^m\text{-ker}(\{y_\sigma\}) \wedge \mathcal{B}_{y_\sigma} = 0_X$  and  $\alpha^m\text{-ker}(\{x_\lambda\}) \leq \mathcal{B}_{y_\sigma}$ ,  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \mathcal{A}_{y_\sigma} = 0_X$  [since  $(X, \tau)$  is a  $F\alpha^m$ - $R_1$ -space and by theorem (4.10)], let  $\delta = \bigwedge \{\mathcal{B}_{y_\sigma} : x_\lambda \in \mathcal{B}_{y_\sigma}\}$ , so we have  $\delta \wedge \mathcal{F} = 0_X$ . Hence  $(X, \tau)$  is a  $F\alpha^m$ - $n$ -space, then there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\mathcal{F} \leq \mathcal{U}$  and  $x_\lambda \in \delta \leq \mathcal{V}$ . Thus,  $(X, \tau)$  is a  $F\alpha^m$ - $r$ -space.

## 5. FUZZY $\alpha^m$ - $T_j$ -SPACES, $j = 0, 1, 2, 3, 4$

**Definition 5.1:** Let  $(X, \tau)$  be a fts. Then  $X$  is called:

- (i) fuzzy  $\alpha^m$ - $T_0$ -space ( $F\alpha^m$ - $T_0$ -space, for short) iff for each pair of distinct fuzzy points in  $X$ , there exists a  $F\alpha^m$ -OS in  $X$  containing one and not the other.
- (ii) fuzzy  $\alpha^m$ - $T_1$ -space ( $F\alpha^m$ - $T_1$ -space, for short) iff for each pair of distinct fuzzy points  $x_\lambda$  and  $y_\sigma$  of  $X$ , there exists  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  containing  $x_\lambda$  and  $y_\sigma$  respectively such that  $y_\sigma \notin \mathcal{U}$  and  $x_\lambda \notin \mathcal{V}$ .
- (iii) fuzzy  $\alpha^m$ - $T_2$ -space ( $F\alpha^m$ - $T_2$ -space, for short) iff for each pair of distinct fuzzy points  $x_\lambda$  and  $y_\sigma$  of  $X$ , there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  in  $X$  such that  $x_\lambda \in \mathcal{U}$  and  $y_\sigma \in \mathcal{V}$ .
- (iv) fuzzy  $\alpha^m$ - $T_3$ -space ( $F\alpha^m$ - $T_3$ -space, for short) iff it is  $F\alpha^m$ - $T_1$ -space and  $F\alpha^m$ - $r$ -space.
- (v) fuzzy  $\alpha^m$ - $T_4$ -space ( $F\alpha^m$ - $T_4$ -space, for short) iff it is  $F\alpha^m$ - $T_1$ -space and  $F\alpha^m$ - $n$ -space.

**Example 5.2:** Let  $X = \{a, b\}$  and  $\tau = \{0_X, a_1, 1_X\}$  be a fts on  $X$ . Then  $a_1$  is a crisp point in  $X$  and  $(X, \tau)$  is a  $F\alpha^m$ - $T_0$ -space.

**Example 5.3:** Let  $X = \{u, v\}$  and  $\tau = \{0_X, u_1, v_1, 1_X\}$  be a fts on  $X$ . Then  $u_1, v_1$  are crisp points in  $X$  and  $(X, \tau)$  is a  $F\alpha^m$ - $T_1$ -space and  $F\alpha^m$ - $T_2$ -space.

**Example 5.4:** The discrete fuzzy topology in  $X = [-2, 2]$  is a  $F\alpha^m$ - $T_3$ -space and  $F\alpha^m$ - $T_4$ -space.

**Remark 5.5:** Every  $F\alpha^m$ - $T_k$ -space is a  $F\alpha^m$ - $T_{k-1}$ -space,  $k = 1, 2, 3, 4$ .

**Proof:** Clearly.

**Theorem 5.6:** A fts  $(X, \tau)$  is  $F\alpha^m$ - $T_0$ -space iff either  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$  or  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$ , for each  $x \neq y \in X$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m$ - $T_0$ -space then for each  $x \neq y \in X$ , there exists a  $F\alpha^m$ -OS  $\mathcal{U}$  such that  $x_\lambda \in \mathcal{U}$ ,  $y_\sigma \notin \mathcal{U}$  or  $x_\lambda \notin \mathcal{U}$ ,  $y_\sigma \in \mathcal{U}$ . Thus either  $x_\lambda \in \mathcal{U}$ ,  $y_\sigma \notin \mathcal{U}$  implies  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$  or  $x_\lambda \notin \mathcal{U}$ ,  $y_\sigma \in \mathcal{U}$  implies  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$ . Conversely, let either  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$  or  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$ , for each  $x \neq y \in X$ . Then there exists a  $F\alpha^m$ -OS  $\mathcal{U}$  such that  $x_\lambda \in \mathcal{U}$ ,  $y_\sigma \notin \mathcal{U}$  or  $x_\lambda \notin \mathcal{U}$ ,  $y_\sigma \in \mathcal{U}$ . Thus  $(X, \tau)$  is a  $F\alpha^m$ - $T_0$ -space.



**Theorem 5.7:** A fts  $(X, \tau)$  is  $F\alpha^m-T_0$ -space iff either  $\alpha^m\text{-ker}(\{x_\lambda\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{y_\sigma\}$  or  $\alpha^m\text{-ker}(\{y_\sigma\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{x_\lambda\}$  for each  $x \neq y \in X$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-T_0$ -space then for each  $x \neq y \in X$ , there exists a  $F\alpha^m$ -OS  $\mathcal{U}$  such that  $x_\lambda \in \mathcal{U}, y_\sigma \notin \mathcal{U}$  or  $x_\lambda \notin \mathcal{U}, y_\sigma \in \mathcal{U}$ . Now if  $x_\lambda \in \mathcal{U}, y_\sigma \notin \mathcal{U}$  implies  $\alpha^m\text{-ker}(\{x_\lambda\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{y_\sigma\}$ . Or if  $x_\lambda \notin \mathcal{U}, y_\sigma \in \mathcal{U}$  implies  $\alpha^m\text{-ker}(\{y_\sigma\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{x_\lambda\}$ .

Conversely, let either  $\alpha^m\text{-ker}(\{x_\lambda\})$  be weakly ultra fuzzy  $\alpha^m$ -separated from  $\{y_\sigma\}$  or  $\alpha^m\text{-ker}(\{y_\sigma\})$  be weakly ultra fuzzy  $\alpha^m$ -separated from  $\{x_\lambda\}$ . Then there exists a  $F\alpha^m$ -OS  $\mathcal{U}$  such that  $\alpha^m\text{-ker}(\{x_\lambda\}) \leq \mathcal{U}$  and  $y_\sigma \notin \mathcal{U}$  or  $\alpha^m\text{-ker}(\{y_\sigma\}) \leq \mathcal{U}, x_\lambda \notin \mathcal{U}$  implies  $x_\lambda \in \mathcal{U}, y_\sigma \notin \mathcal{U}$  or  $x_\lambda \notin \mathcal{U}, y_\sigma \in \mathcal{U}$ . Thus,  $(X, \tau)$  is a  $F\alpha^m-T_0$ -space.

**Theorem 5.8:** A fts  $(X, \tau)$  is  $F\alpha^m-T_1$ -space iff for each  $x \neq y \in X$ ,  $\alpha^m\text{-ker}(\{x_\lambda\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{y_\sigma\}$  and  $\alpha^m\text{-ker}(\{y_\sigma\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{x_\lambda\}$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-T_1$ -space then for each  $x \neq y \in X$ , there exist  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $x_\lambda \in \mathcal{U}, y_\sigma \notin \mathcal{U}$  and  $x_\lambda \notin \mathcal{V}, y_\sigma \in \mathcal{V}$ . Implies  $\alpha^m\text{-ker}(\{x_\lambda\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{y_\sigma\}$  and  $\alpha^m\text{-ker}(\{y_\sigma\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{x_\lambda\}$ .

Conversely, let  $\alpha^m\text{-ker}(\{x_\lambda\})$  be weakly ultra fuzzy  $\alpha^m$ -separated from  $\{y_\sigma\}$  and  $\alpha^m\text{-ker}(\{y_\sigma\})$  be weakly ultra fuzzy  $\alpha^m$ -separated from  $\{x_\lambda\}$ . Then there exist  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\alpha^m\text{-ker}(\{x_\lambda\}) \leq \mathcal{U}, y_\sigma \notin \mathcal{U}$  and  $\alpha^m\text{-ker}(\{y_\sigma\}) \leq \mathcal{V}, x_\lambda \notin \mathcal{V}$  implies  $x_\lambda \in \mathcal{U}, y_\sigma \notin \mathcal{U}$  and  $x_\lambda \notin \mathcal{V}, y_\sigma \in \mathcal{V}$ . Thus,  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space.

**Theorem 5.9:** A fts  $(X, \tau)$  is  $F\alpha^m-T_1$ -space iff for each  $x \in X$ ,  $\alpha^m\text{-ker}(\{x_\lambda\}) = \{x_\lambda\}$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-T_1$ -space and let  $\alpha^m\text{-ker}(\{x_\lambda\}) \neq \{x_\lambda\}$ . Then  $\alpha^m\text{-ker}(\{x_\lambda\})$  contains another fuzzy point distinct from  $x_\lambda$  say  $y_\sigma$ . So  $y_\sigma \in \alpha^m\text{-ker}(\{x_\lambda\})$  implies  $\alpha^m\text{-ker}(\{x_\lambda\})$  is not weakly ultra fuzzy  $\alpha^m$ -separated from  $\{y_\sigma\}$ . Hence by theorem (5.8),  $(X, \tau)$  is not a  $F\alpha^m-T_1$ -space this is contradiction. Thus  $\alpha^m\text{-ker}(\{x_\lambda\}) = \{x_\lambda\}$ .

Conversely, let  $\alpha^m\text{-ker}(\{x_\lambda\}) = \{x_\lambda\}$ , for each  $x \in X$  and let  $(X, \tau)$  be not a  $F\alpha^m-T_1$ -space. Then by theorem (5.8),  $\alpha^m\text{-ker}(\{x_\lambda\})$  is not weakly ultra fuzzy  $\alpha^m$ -separated from  $\{y_\sigma\}$ , this means that for every  $F\alpha^m$ -OS  $\mathcal{U}$  contains  $\alpha^m\text{-ker}(\{x_\lambda\})$  then  $y_\sigma \in \mathcal{U}$  implies  $y_\sigma \in \bigwedge \{\mathcal{U} \in F\alpha^m-O(X) : x_\lambda \in \mathcal{U}\}$  implies  $y_\sigma \in \alpha^m\text{-ker}(\{x_\lambda\})$ , this is contradiction. Thus,  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space.

**Theorem 5.10:** A fts  $(X, \tau)$  is  $F\alpha^m-T_1$ -space iff for each  $x \neq y \in X$ ,  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$  and  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-T_1$ -space then for each  $x \neq y \in X$ , there exists  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $x_\lambda \in \mathcal{U}, y_\sigma \notin \mathcal{U}$  and  $y_\sigma \in \mathcal{V}, x_\lambda \notin \mathcal{V}$ . Implies  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$  and  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$ .

Conversely, let  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$  and  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$ , for each  $x \neq y \in X$ . Then there exists  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $x_\lambda \in \mathcal{U}, y_\sigma \notin \mathcal{U}$  and  $y_\sigma \in \mathcal{V}, x_\lambda \notin \mathcal{V}$ . Thus,  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space.

**Theorem 5.11:** A fts  $(X, \tau)$  is  $F\alpha^m-T_1$ -space iff for each  $x \neq y \in X$  implies  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \alpha^m\text{-ker}(\{y_\sigma\}) = 0_X$ .

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-T_1$ -space. Then  $\alpha^m\text{-ker}(\{x_\lambda\}) = \{x_\lambda\}$  and  $\alpha^m\text{-ker}(\{y_\sigma\}) = \{y_\sigma\}$  [by theorem (5.9)]. Thus,  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \alpha^m\text{-ker}(\{y_\sigma\}) = 0_X$ .

Conversely, let for each  $x \neq y \in X$  implies  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \alpha^m\text{-ker}(\{y_\sigma\}) = 0_X$  and let  $(X, \tau)$  be not a  $F\alpha^m-T_1$ -space then for each  $x \neq y \in X$  implies  $y_\sigma \in \alpha^m\text{-ker}(\{x_\lambda\})$  or  $x_\lambda \in \alpha^m\text{-ker}(\{y_\sigma\})$  [by theorem (5.10)], then  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \alpha^m\text{-ker}(\{y_\sigma\}) \neq 0_X$  this is contradiction. Thus,  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space.

**Theorem 5.12:** A fts  $(X, \tau)$  is  $F\alpha^m-T_1$ -space iff  $(X, \tau)$  is  $F\alpha^m-T_0$ -space and  $F\alpha^m-R_0$ -space.

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-T_1$ -space and let  $x_\lambda \in \mathcal{U}$  be a  $F\alpha^m$ -OS, then for each  $x \neq y \in X$ ,  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \alpha^m\text{-ker}(\{y_\sigma\}) = 0_X$  [by theorem (5.11)] implies  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$  and  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$  this means  $\alpha^m\text{-cl}(\{x_\lambda\}) = \{x_\lambda\}$ , hence  $\alpha^m\text{-cl}(\{x_\lambda\}) \leq \mathcal{U}$ . Thus,  $(X, \tau)$  is a  $F\alpha^m-R_0$ -space.

Conversely, let  $(X, \tau)$  be a  $F\alpha^m-T_0$ -space and  $F\alpha^m-R_0$ -space, then for each  $x \neq y \in X$  there exists a  $F\alpha^m$ -OS  $\mathcal{U}$  such that  $x_\lambda \in \mathcal{U}$ ,  $y_\sigma \notin \mathcal{U}$  or  $x_\lambda \notin \mathcal{U}$ ,  $y_\sigma \in \mathcal{U}$ . Say  $x_\lambda \in \mathcal{U}$ ,  $y_\sigma \notin \mathcal{U}$  since  $(X, \tau)$  is a  $F\alpha^m-R_0$ -space, then  $\alpha^m-cl(\{x_\lambda\}) \leq \mathcal{U}$ , this means there exists a  $F\alpha^m$ -OS  $\mathcal{V}$  such that  $y_\sigma \in \mathcal{V}$ ,  $x_\lambda \notin \mathcal{V}$ . Thus,  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space.

**Theorem 5.13:** A fts  $(X, \tau)$  is  $F\alpha^m-T_2$ -space iff

- (i)  $(X, \tau)$  is  $F\alpha^m-T_0$ -space and  $F\alpha^m-R_1$ -space.
- (ii)  $(X, \tau)$  is  $F\alpha^m-T_1$ -space and  $F\alpha^m-R_1$ -space.

**Proof:** (i) Let  $(X, \tau)$  be a  $F\alpha^m-T_2$ -space then it is a  $F\alpha^m-T_0$ -space. Now since  $(X, \tau)$  is a  $F\alpha^m-T_2$ -space then for each  $x \neq y \in X$ , there exist disjoint  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $x_\lambda \in \mathcal{U}$  and  $y_\sigma \in \mathcal{V}$  implies  $x_\lambda \notin \alpha^m-cl(\{y_\sigma\})$  and  $y_\sigma \notin \alpha^m-cl(\{x_\lambda\})$ , therefore  $\alpha^m-cl(\{x_\lambda\}) = \{x_\lambda\} \leq \mathcal{U}$  and  $\alpha^m-cl(\{y_\sigma\}) = \{y_\sigma\} \leq \mathcal{V}$ . Thus,  $(X, \tau)$  is a  $F\alpha^m-R_1$ -space.

Conversely, let  $(X, \tau)$  be a  $F\alpha^m-T_0$ -space and  $F\alpha^m-R_1$ -space, then for each  $x \neq y \in X$ , there exists a  $F\alpha^m$ -OS  $\mathcal{U}$  such that  $x_\lambda \in \mathcal{U}$ ,  $y_\sigma \notin \mathcal{U}$  or  $y_\sigma \in \mathcal{U}$ ,  $x_\lambda \notin \mathcal{U}$ , implies  $\alpha^m-cl(\{x_\lambda\}) \neq \alpha^m-cl(\{y_\sigma\})$ , since  $(X, \tau)$  is a  $F\alpha^m-R_1$ -space [by assumption], then there exist disjoint  $F\alpha^m$ -OS  $\mathcal{M}, \mathcal{N}$  such that  $x_\lambda \in \mathcal{M}$  and  $y_\sigma \in \mathcal{N}$ . Thus,  $(X, \tau)$  is a  $F\alpha^m-T_2$ -space.

(ii) By the same way of part (i) a  $F\alpha^m-T_2$ -space is  $F\alpha^m-T_1$ -space and  $F\alpha^m-R_1$ -space.

Conversely, let  $(X, \tau)$  be a  $F\alpha^m-T_1$ -space and  $F\alpha^m-R_1$ -space, then for each  $x \neq y \in X$ , there exist  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $x_\lambda \in \mathcal{U}$ ,  $y_\sigma \notin \mathcal{U}$  and  $y_\sigma \in \mathcal{V}$ ,  $x_\lambda \notin \mathcal{V}$  implies  $\alpha^m-cl(\{x_\lambda\}) \neq \alpha^m-cl(\{y_\sigma\})$ , since  $(X, \tau)$  is a  $F\alpha^m-R_1$ -space, then there exist disjoint  $F\alpha^m$ -OS  $\mathcal{M}, \mathcal{N}$  such that  $x_\lambda \in \mathcal{M}$  and  $y_\sigma \in \mathcal{N}$ . Thus,  $(X, \tau)$  is a  $F\alpha^m-T_2$ -space.

**Corollary 5.14:** A  $F\alpha^m-T_0$ -space is  $F\alpha^m-T_2$ -space iff for each  $x \neq y \in X$  with  $\alpha^m-ker(\{x_\lambda\}) \neq \alpha^m-ker(\{y_\sigma\})$  then there exist  $F\alpha^m$ -CS  $\mathcal{A}_1, \mathcal{A}_2$  such that  $\alpha^m-ker(\{x_\lambda\}) \leq \mathcal{A}_1$ ,  $\alpha^m-ker(\{x_\lambda\}) \wedge \mathcal{A}_2 = 0_X$  and  $\alpha^m-ker(\{y_\sigma\}) \leq \mathcal{A}_2$ ,  $\alpha^m-ker(\{y_\sigma\}) \wedge \mathcal{A}_1 = 0_X$  and  $\mathcal{A}_1 \vee \mathcal{A}_2 = 1_X$ .

**Proof:** By theorem (4.10) and theorem (5.13).

**Corollary 5.15:** A  $F\alpha^m-T_1$ -space is  $F\alpha^m-T_2$ -space iff one of the following conditions holds:

- (i) for each  $x \neq y \in X$  with  $\alpha^m-cl(\{x_\lambda\}) \neq \alpha^m-cl(\{y_\sigma\})$ , then there exist  $F\alpha^m$ -OS  $\mathcal{U}, \mathcal{V}$  such that  $\alpha^m-cl(\alpha^m-ker(\{x_\lambda\})) \leq \mathcal{U}$  and  $\alpha^m-cl(\alpha^m-ker(\{y_\sigma\})) \leq \mathcal{V}$ .
- (ii) for each  $x \neq y \in X$  with  $\alpha^m-ker(\{x_\lambda\}) \neq \alpha^m-ker(\{y_\sigma\})$ , then there exist  $F\alpha^m$ -CS  $\mathcal{A}_1, \mathcal{A}_2$  such that  $\alpha^m-ker(\{x_\lambda\}) \leq \mathcal{A}_1$ ,  $\alpha^m-ker(\{x_\lambda\}) \wedge \mathcal{A}_2 = 0_X$  and  $\alpha^m-ker(\{y_\sigma\}) \leq \mathcal{A}_2$ ,  $\alpha^m-ker(\{y_\sigma\}) \wedge \mathcal{A}_1 = 0_X$  and  $\mathcal{A}_1 \vee \mathcal{A}_2 = 1_X$ .

**Proof:** (i) By corollary (4.11) and theorem (5.13).

(ii) By theorem (4.10) and theorem (5.13).

**Theorem 5.16:** A  $F\alpha^m-R_1$ -space is  $F\alpha^m-T_2$ -space iff one of the following conditions holds:

- (i) for each  $x \in X$ ,  $\alpha^m-ker(\{x_\lambda\}) = \{x_\lambda\}$ .
- (ii) for each  $x \neq y \in X$ ,  $\alpha^m-ker(\{x_\lambda\}) \neq \alpha^m-ker(\{y_\sigma\})$  implies  $\alpha^m-ker(\{x_\lambda\}) \wedge \alpha^m-ker(\{y_\sigma\}) = 0_X$ .
- (iii) for each  $x \neq y \in X$ , either  $x_\lambda \notin \alpha^m-ker(\{y_\sigma\})$  or  $y_\sigma \notin \alpha^m-ker(\{x_\lambda\})$ .
- (iv) for each  $x \neq y \in X$ , then  $x_\lambda \notin \alpha^m-ker(\{y_\sigma\})$  and  $y_\sigma \notin \alpha^m-ker(\{x_\lambda\})$ .

**Proof:** (i) Let  $(X, \tau)$  be a  $F\alpha^m-T_2$ -space. Then  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space and  $F\alpha^m-R_1$ -space [by theorem (5.13)]. Hence by theorem (5.9),  $\alpha^m-ker(\{x_\lambda\}) = \{x_\lambda\}$  for each  $x \in X$ .

Conversely, let for each  $x \in X$ ,  $\alpha^m-ker(\{x_\lambda\}) = \{x_\lambda\}$ , then by theorem (5.9),  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space. Also  $(X, \tau)$  is a  $F\alpha^m-R_1$ -space by assumption. Hence by theorem (5.13),  $(X, \tau)$  is a  $F\alpha^m-T_2$ -space.

(ii) Let  $(X, \tau)$  be a  $F\alpha^m-T_2$ -space. Then  $(X, \tau)$  is  $F\alpha^m-T_1$ -space [by remark (5.5)]. Hence by theorem (5.11),  $\alpha^m-ker(\{x_\lambda\}) \wedge \alpha^m-ker(\{y_\sigma\}) = 0_X$  for each  $x \neq y \in X$ .

Conversely, assume that for each  $x \neq y \in X$ ,  $\alpha^m-ker(\{x_\lambda\}) \neq \alpha^m-ker(\{y_\sigma\})$  implies  $\alpha^m-ker(\{x_\lambda\}) \wedge \alpha^m-ker(\{y_\sigma\}) = 0_X$ . So by theorem (5.11),  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space, also  $(X, \tau)$  is a  $F\alpha^m-R_1$ -space by assumption. Hence by theorem (5.13),  $(X, \tau)$  is a  $F\alpha^m-T_2$ -space.

(iii) Let  $(X, \tau)$  be a  $F\alpha^m-T_2$ -space. Then  $(X, \tau)$  is a  $F\alpha^m-T_0$ -space [by remark (5.5)]. Hence by theorem (5.6), either  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$  or  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$  for each  $x \neq y \in X$ .

Conversely, assume that for each  $x \neq y \in X$ , either  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$  or  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$  for each  $x \neq y \in X$ . So by theorem (5.6),  $(X, \tau)$  is a  $F\alpha^m-T_0$ -space, also  $(X, \tau)$  is  $F\alpha^m-R_1$ -space by assumption. Thus  $(X, \tau)$  is a  $F\alpha^m-T_2$ -space [by theorem (5.13)].

(iv) Let  $(X, \tau)$  be a  $F\alpha^m-T_2$ -space. Then  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space and  $F\alpha^m-R_1$ -space [by theorem (5.13)]. Hence by theorem (5.10),  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$  and  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$ .

Conversely, let for each  $x \neq y \in X$  then  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$  and  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$ . Then by theorem (5.10),  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space. Also  $(X, \tau)$  is a  $F\alpha^m-R_1$ -space by assumption. Hence by theorem (5.13),  $(X, \tau)$  is a  $F\alpha^m-T_2$ -space.

**Remark 5.17:** Each fuzzy separation axiom is defined as the conjunction of two weaker axioms:  $F\alpha^m-T_k$ -space =  $F\alpha^m-R_{k-1}$ -space and  $F\alpha^m-T_{k-1}$ -space =  $F\alpha^m-R_{k-1}$ -space and  $F\alpha^m-T_0$ -space,  $k = 1, 2, 3, 4$ .

**Theorem 5.18:** Let  $(X, \tau)$  be a fts and  $\alpha^m\text{-ker}(\{x_\lambda\}) = \{x_\lambda\}$  for each  $x \in X$  then  $(X, \tau)$  is  $F\alpha^m-T_3$ -space if and only if it is a  $F\alpha^m-R_2$ -space.

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-T_3$ -space. Then,  $(X, \tau)$  is a  $F\alpha^m-R_2$ -space [By remark (5.17)].

Conversely, let  $(X, \tau)$  be a  $F\alpha^m-R_2$ -space then it is a  $F\alpha^m r$ -space [definition (4.12)(iii)]. By assumption,  $\alpha^m\text{-ker}(\{x_\lambda\}) = \{x_\lambda\}$  for each  $x \in X$ , then  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space [by theorem (5.9)]. Hence by remark (5.17),  $(X, \tau)$  is a  $F\alpha^m-T_3$ -space.

**Theorem 5.19:** Let  $(X, \tau)$  be a fts and let  $x \neq y \in X$ , implies  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \alpha^m\text{-ker}(\{y_\sigma\}) = 0_X$ , then  $(X, \tau)$  is a  $F\alpha^m-T_3$ -space iff it is a  $F\alpha^m-R_2$ -space.

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-T_3$ -space. Then  $(X, \tau)$  is a  $F\alpha^m-R_2$ -space [by remark (5.17)].

Conversely, let  $(X, \tau)$  be a  $F\alpha^m-R_2$ -space then it is a  $F\alpha^m r$ -space [definition (4.12)(iii)]. By assumption,  $\alpha^m\text{-ker}(\{x_\lambda\}) \wedge \alpha^m\text{-ker}(\{y_\sigma\}) = 0_X$ , for each  $x \neq y \in X$ , then by theorem (5.11),  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space. Hence by remark (5.17),  $(X, \tau)$  is a  $F\alpha^m-T_3$ -space.

**Theorem 5.20:** Let  $(X, \tau)$  be a fts and for each  $x \neq y \in X$  either  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$  or  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$ , then  $(X, \tau)$  is a  $F\alpha^m-T_3$ -space iff it is a  $F\alpha^m-R_2$ -space.

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-T_3$ -space. Then  $(X, \tau)$  is a  $F\alpha^m-R_2$ -space [by remark (5.17)].

Conversely, let  $(X, \tau)$  be a  $F\alpha^m-R_2$ -space then it is a  $F\alpha^m r$ -space [definition (4.12)(iii)]. By assumption, for each  $x \neq y \in X$  either  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$  or  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$ . This means either  $\alpha^m\text{-ker}(\{x_\lambda\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{y_\sigma\}$  or  $\alpha^m\text{-ker}(\{y_\sigma\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{x_\lambda\}$ , so by theorem (5.7),  $(X, \tau)$  is a  $F\alpha^m-T_0$ -space. Hence by remark (5.17),  $(X, \tau)$  is a  $F\alpha^m-T_3$ -space.

**Theorem 5.21:** Let  $(X, \tau)$  be a fts and let  $x \neq y \in X$ , then  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$  and  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$ ,  $(X, \tau)$  is a  $F\alpha^m-T_3$ -space iff it is a  $F\alpha^m-R_2$ -space.

**Proof:** Let  $(X, \tau)$  be a  $F\alpha^m-T_3$ -space. Then  $(X, \tau)$  is a  $F\alpha^m-R_2$ -space [by remark (5.17)].

Conversely, let  $(X, \tau)$  be a  $F\alpha^m-R_2$ -space then it is a  $F\alpha^m r$ -space [definition (4.12)(iii)]. By assumption, for each  $x \neq y \in X$  then  $x_\lambda \notin \alpha^m\text{-ker}(\{y_\sigma\})$  and  $y_\sigma \notin \alpha^m\text{-ker}(\{x_\lambda\})$ . Therefore,  $\alpha^m\text{-ker}(\{x_\lambda\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{y_\sigma\}$  and  $\alpha^m\text{-ker}(\{y_\sigma\})$  is weakly ultra fuzzy  $\alpha^m$ -separated from  $\{x_\lambda\}$ , so by theorem (5.8),  $(X, \tau)$  is a  $F\alpha^m-T_1$ -space. Hence by remark (5.17),  $(X, \tau)$  is a  $F\alpha^m-T_3$ -space.

**Remark 5.22:** The relation between fuzzy  $\alpha^m$ -separation axioms can be representing as a matrix. Therefore, the element  $a_{ij}$  refers to this relation. As the following matrix representation shows:

and	$F\alpha^m-T_0$	$F\alpha^m-T_1$	$F\alpha^m-T_2$	$F\alpha^m-T_3$	$F\alpha^m-T_4$	$F\alpha^m-R_0$	$F\alpha^m-R_1$	$F\alpha^m-R_2$	$F\alpha^m-R_3$
$F\alpha^m-T_0$	$F\alpha^m-T_0$	$F\alpha^m-T_1$	$F\alpha^m-T_2$	$F\alpha^m-T_3$	$F\alpha^m-T_4$	$F\alpha^m-T_1$	$F\alpha^m-T_2$	$F\alpha^m-R_3$	$F\alpha^m-T_4$
$F\alpha^m-T_1$	$F\alpha^m-T_1$	$F\alpha^m-T_1$	$F\alpha^m-T_2$	$F\alpha^m-T_3$	$F\alpha^m-T_4$	$F\alpha^m-T_1$	$F\alpha^m-T_2$	$F\alpha^m-R_3$	$F\alpha^m-T_4$
$F\alpha^m-T_2$	$F\alpha^m-T_2$	$F\alpha^m-T_2$	$F\alpha^m-T_2$	$F\alpha^m-T_3$	$F\alpha^m-T_4$	$F\alpha^m-T_2$	$F\alpha^m-T_2$	$F\alpha^m-R_3$	$F\alpha^m-T_4$
$F\alpha^m-T_3$	$F\alpha^m-T_3$	$F\alpha^m-T_3$	$F\alpha^m-T_3$	$F\alpha^m-T_3$	$F\alpha^m-T_4$	$F\alpha^m-T_3$	$F\alpha^m-T_3$	$F\alpha^m-T_3$	$F\alpha^m-T_4$
$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-T_4$
$F\alpha^m-R_0$	$F\alpha^m-T_1$	$F\alpha^m-T_1$	$F\alpha^m-T_2$	$F\alpha^m-T_3$	$F\alpha^m-T_4$	$F\alpha^m-R_0$	$F\alpha^m-R_1$	$F\alpha^m-R_2$	$F\alpha^m-R_3$
$F\alpha^m-R_1$	$F\alpha^m-T_2$	$F\alpha^m-T_2$	$F\alpha^m-T_2$	$F\alpha^m-T_3$	$F\alpha^m-T_4$	$F\alpha^m-R_1$	$F\alpha^m-R_1$	$F\alpha^m-R_2$	$F\alpha^m-R_3$
$F\alpha^m-R_2$	$F\alpha^m-T_3$	$F\alpha^m-T_3$	$F\alpha^m-T_3$	$F\alpha^m-T_3$	$F\alpha^m-T_4$	$F\alpha^m-R_2$	$F\alpha^m-R_2$	$F\alpha^m-R_2$	$F\alpha^m-R_3$
$F\alpha^m-R_3$	$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-T_4$	$F\alpha^m-R_3$	$F\alpha^m-R_3$	$F\alpha^m-R_3$	$F\alpha^m-R_3$

#### Matrix Representation

The relation between fuzzy  $\alpha^m$ -separation axioms

#### REFERENCES

- [1] S. Bin Shahna, "On fuzzy strong semi continuity and fuzzy pre continuity", *Fuzzy sets and system*, 44(2), (1991), 303-308.
- [2] Sikn, "On fuzzy FC - Compactness", *Comm. Korean Math. Soc.*, 13(1998), 137-150.
- [3] L. Chang, "Fuzzy topological spaces", *J. Math. Anal. Appl.*, 24(1968), 182-190.
- [4] K. Wong, "Fuzzy points and local properties of fuzzy topology", *J. Math. Anal. Appl.*, 46(1974), 316-328.
- [5] G. Balasubramania and P. Sundaram, "On some generalizations of fuzzy continuous functions", *Fuzzy sets and Systems*, 86(1), (1997), 93-100.
- [6] L. A. Zadeh, "Fuzzy sets", *Information and control*, 8 (1965), 338-353.
- [7] M. Mathew and R. Parimelazhagan, " $\alpha^m$ -Closed set in topological spaces", *International Journal of Mathematical Analysis*, Vol.8, No.47, (2014), 2325-2329.
- [8] P. M. Pu and Y. M. Liu, "Fuzzy topology I. Neighborhood structure of a fuzzy point and Moore-Smith convergence", *J. Math. Anal. Appl.*, 76(1980), 571-599.
- [9] R. K. Saraf and S. Mishra, "Fg $\alpha$ -closed sets", *J. Tripura Math. Soc.*, 2(2000), 27-32.
- [10] R. K. Saraf, G. Navalagi and M. Khanna, "On fuzzy semi pre generalized closed sets", *Bull. Malaysian Math. Sci. Soc.*, 28(1), (2005), 19-30.
- [11] S. S. Thakur and S. Singh, "On fuzzy semi pre open sets and fuzzy semi pre continuous", *Fuzzy sets and system*, 98(3), (1998), 383-391.
- [12] S. Ganguly and S. Saha, "A note on  $\delta$ -continuity and  $\delta$ -connected sets in fuzzy set theory", *Simon Stevin*, 62(1988).